

ELECTRONIC SCORE PREDICTIVE OF ENDOMETRIOSIS

ENDOMETRIOSIS INDEX: A SOFTWARE-DERIVED SCORE TO PREDICT THE PRESENCE AND SEVERITY OF THE DISEASE

Alessandro Fasciani, M.D.,^a Salvo Reina,^b Mauro Costa, M.D.,^c Matteo Puntoni, Ph.D.,^d
Guido Bocci, M.D. Ph.D.,^e Felice Repetti, M.D.^a

^a Division of Gynecology and Obstetrics - Department of Genetic, Perinatal and Gynecological Sciences - Galliera Hospital - Genoa - Italy.

^b Laboratory of experimental microbiology and epidemiology - Department Discat - School of Medicine - University of Genoa - Italy.

^c Infertility Center, Department of Genetic, Perinatal and Gynecological Sciences - Galliera Hospital - Genoa - Italy.

^d Medical Oncology and Clinical Trial Unit - Galliera Hospital - Genoa – Italy

^e Division of Pharmacology and Chemotherapy - Department of Internal Medicine – University of Pisa – Pisa - Italy

Reprint requests:

Alessandro Fasciani, M.D.,

Division of Gynecology and Obstetrics – Mura delle Cappuccine, 14 - Galliera Hospital – 16124 Genoa - Italy.

Tel: ++39 010 5634945 – Fax: ++39 010 5634705 E-mail: alessandro.fasciani@galliera.it

'Endometriosis Index' has been successfully tested by physicians as a non-invasive tool to score the presence of endometriosis in patients with suspected disease.

ABSTRACT

Objective: To develop a clinical index that would positively predict the presence of endometriosis.

Design: Prospective single-centre observational study.

Setting: A hospital-based institute.

Patient(s): 120 patients affected by chronic pelvic pain, infertility or with clinical suspicion of endometriosis.

Intervention(s): Electronic processing of clinical data with software-assistance at the end of each consultation and digital video recording of surgeries.

Main Outcome Measure(s): Endometriosis Index (EI), the score calculated using clinical parameters correlated with macroscopic/microscopic presence or absence of endometriosis.

Result(s): Endometriosis was staged and treated in 95 cases, the remaining 25 women presented benign pathology with no endometriosis. Patients with positive operative findings of endometriosis had a mean (\pm standard deviation) pre-operative EI score of 22 ± 12 , while mean EI value of patients with no operative findings of endometriosis was 8 ± 6 ($p < 0.001$). Cumulative distribution of EI shows increasing values from controls to peritoneal, ovarian and deeply infiltrating endometriosis (DIE) (p for trend < 0.001). A Logistic model showed an OR = 24.7 (95%CI = 8.3 to 73.7) to have DIE for women with EI score > 28 (75th percentile value) versus women with EI score ≤ 28 .

Conclusion(s): These data suggest that a dedicated Endometriosis Index is effective in identifying patients who would benefit from early surgical management. We propose the use of this non-invasive tool to reduce the delay between the onset of symptoms and a surgical diagnosis of endometriosis.

Key Words: Endometriosis, Index, Software, Diagnosis.

INTRODUCTION

Endometriosis has significant medical and social impact world-wide. The overall delay between onset of symptoms and actual diagnosis has been calculated 9.28 years (1). Pooled data from a 9-year single-centre study have estimated the prevalence of endometriosis between 8 and 10% (2). At an individual level, chronic pelvic pain leads to years of disability and suffering, with loss of employment, marital discord, divorce and numerous untoward and unsuccessful medical misadventures (3). Clearly, pelvic pain is an important issue in the health care of women contributing to 10% of all outpatient gynaecological visits, is responsible for approximately 40% of laparoscopies, and is the indication for 10% - 15% of hysterectomies (4). The existence of a relationship between chronic pelvic pain symptoms and endometriosis is widely accepted, but various painful pelvic symptoms are normally present in the general population (5). The complex nature of chronic pain and our naïve understanding of its origin, particularly in the setting of endometriosis, results in the use of therapies that generally provide only temporary relief of symptoms (6). The American College and the Royal College of Obstetricians and Gynaecologists have both recently recommend the empirical use of medical therapy before confirming a definitive diagnosis when the risks and benefits of empiric surgery are considered (7). However, endometriosis may be progressive and early laparoscopic diagnosis in patients suffering from this potentially serious disease could be appropriate. Recognizing the known effect and effectiveness of surgical excision on endometriosis (8), the dilemma for individual clinicians is to choose when to operate. Despite several classification systems and questionnaires developed thus far, no standard reference exists for patients suffering from this disabling condition. We propose a tool to determine, at the time of the first consultation, whether a woman would benefit from early surgery to treat her condition versus simply empirical treatment. To address this issue we have created a non-invasive diagnostic Endometriosis Index (EI) from 38 variables and

parameters derived by the patient pain evaluation, physician consultation, and diagnostic evidence.

We propose creating a standardized index, taking into consideration evidence-based signs, symptoms and diagnostic criteria (9-13), to preoperatively determine the patients probability of having endometriosis and determining which patients will gain the greatest benefit form early surgery.

MATERIALS AND METHODS

Population

120 patients, referred to our unit for suspicion of endometriosis, chronic pelvic pain and infertility, were randomly and prospectively evaluated with software-assistance before and after surgery. All women were operated and videos were recorded via hospital intranet. At the time of surgery endometriosis was staged according to the Revised American Society for Reproductive Medicine classification (14) and each case was confirmed by histology. On the base of the information received from surgeons and pathologists endometriosis was sub-classified in peritoneal forms (endometriotic extension < 1 mm underneath the peritoneum), ovarian endometriomata (intra-ovarian endometriotic lesions) and deeply infiltrating endometriosis (DIE) (endometriotic extension > 5 mm underneath the peritoneum) (15, 16). The study was reviewed and approved by the local Institutional Review Board.

Model description

The unified evaluation model is a software real-time calculation based on homothetic transformation (17) of 38 variables and parameters expressed with a normalised score. The exhaustiveness of the symptoms studied was ensured by basing the list on a comprehensive literature review (1, 3-13, 15, 18-22) and on our clinical practice. Endometriosis health profile questionnaires, health-related quality of life instruments and pain evaluation systems previously used in clinical studies (3, 8-10, 13, 18, 23-30) have been carefully evaluated and constitute the educational base of this software assisted health profile assessment.

The final score of EI is generated to be the quantification of the pathological status at the time of each consultation. The variables and parameters were distributed over three components: i) health profile assessment of the patient, ii) medical consultation and iii) diagnostic evidences.

This design is adherent with the model structure and the foreseen questionnaire can be filled out on paper or entered directly on the computer. Each component has a collection of values (indicators) grouped according to a logical cascade of questions in the following sections:

- i-a) Pain assessment is the result of the following subitems: *1 Dysmenorrhea (IDM) + 2 Non menstrual pelvic pain (IDP) + 3 Dyspareunia (DRS)* (Fig.1A); Characterization of the sensory dimension of pain (CDD) is the result of the following subitems: *4 Throbbing + 5 Stabbing + 6 Cramping + 7 Hot-burning + 8 Heavy + 9 Tender* (Fig.1B).
- i-b) Quality of life limitation (LDD) is the result of the following subitems: *10 Work/School days of absence + 11 Daily activity restriction and 12 Sleep impairment* (Fig.1B),
- i-c: Induced dysfunction and physical alteration (SPA) is the result of the following subitems: gastrointestinal symptoms (*13 Alternating constipation and diarrhoea + 14 Rectal tenesmus + 15 constipation + 16 diarrhoea + 17 Rectal pain*), urinary symptoms (*18 Urinary pain – 19 bladder tenesmus – 20 Frequent urination – 21 Dysuria*) and headache (*22 Headache – 23 Migraine – 24 cluster headache*) (Fig.1C).
- ii + iii: Pelvic examination (VM) is the result of the following subitems: (*25 Grade of fixed uterus + 26 Presence of adnexal cyst(s) + 27 Grade of Douglas tenderness + 28 Presence of infiltrating nodule(s) + 29 Grade of pain at digital exploration*); Diagnostic evidences (IS) result of the following subitems: (*30 CA-125 blood assay + 31 Ultrasound finding(s) + 32 CA 19.9 blood assay + 33 Magnetic resonance finding(s) + 34 X-Ray / Computerized Tomography finding(s) + 35 Colonoscopy finding(s)*) (Fig.1D).

Each record profile is compiled through a sequential data-entry of four panels (Fig.1) and the algorithm calculates EI and plots a nomogram on real-time. We present the original Italian language panels as entered by our patient cohort. Other language versions will be available

soon. All indicators are expressed as visual analogue scales (VAS), the left and the right extremes represent respectively the lowest and the maximum value; each variable or parameter contributes to the final EI score on the basis of its weight that it has been assigned to reduce or emphasise its relevance. In the first phase of use, the physicians can over-ride the default setting of the software on the basis of their clinical experience. The operator has the possibility to easily adjust the indicator's weighting

An inherent model feature, the algorithm calculates meaningful EI values, even in the case of missing or incomplete information. In fact, in real practise, it is unlikely to cover all the data-entry panels when interviewing patients. Beside logistic reasons or practical limitations for having incomplete information, the model also accommodates these critical events and allows for data validation. For instance, it is not possible to consider the indicator score for sexual intercourse pain when, say, a young patient had not (a case where it would not make sense to consider a zero score for this type of pain, because there were no sexual intercourses). The algorithm considers the relative weighting of an indicator (either variable or parameter) by discriminating between zero and *nil* values, so distinguishing a zero scalar number from the absence of indicator (when not applicable or not meaningful). In this latter case, the redistribution of the algebraic, cumulative sum of indicators will only take into account the meaningful values and the EI will be calculated accordingly.

The information hierarchy is therefore treated by the model so that the EI values can still be calculated even when entire sections are not entered. EI software considers several approximation possibilities and allows the physician to decide whether to exclude or emphasise specific information. For example each symptom reported in the field i-c (Fig.1C) makes the EI higher when correlated to the menses; each of these parameters, in fact, is classified as catamenial by clicking on the check-box named "*synchronous with cycle*".

Several patients can be followed over time so that an auto-correlation and a trends analysis give an epidemiological prospect. The software provides a storage template of the heuristic weighted knowledge for further uses.

Statistical Analysis

Categorical data were summarized as number (percentage) of subjects while continuous data as mean \pm standard deviation (SD) or as median (minimum – maximum). Univariate associations between endometriosis status and all other variables considered were assessed using chi-square or Fisher exact test analyses for categorical variables and the Kruskal-Wallis, Wilcoxon and Mann Whitney tests (in case of non-normal distributions) for continuous variables. Boxplots were used to study distributions of EI score and ASRM score in controls women and in women with different types of endometriosis. A non parametric test for trend across ordered groups was used. Pearson correlation coefficient was used to assess correlation between continuous variables.

A logistic regression model was used to estimate the association between DIE disease and EI score percentiles, adjusting for age of women; the likelihood ratio test was used to assess the statistical significance. Two-tailed probabilities were reported and the p-value of 0.05 was used to define nominal statistical significance. All analyses were conducted using Stata (version 10; StataCorp., College Station, TX) software.

RESULTS

Endometriosis was staged and treated in 95 cases (the remaining 25 women presented benign pathology with no endometriosis nor adhesions). Clinical characteristics of patients are shown in table 1.

The software EI calculated before surgery in patients with proven endometriosis resulted in significantly higher values than those of women with no endometriosis (22.3 ± 11.6 vs 8.1 ± 6.1 , $p < 0.001$). The EI scores obtained at the consultations after the excision of endometriosis (25 ± 15 days) become statistically lower than the values recorded in the same patients before surgery (3.9 ± 2.5 vs 22.3 ± 11.6 , $p < 0.001$) and similar to EI calculated in the group of control (Table 1) (Figure 2).

Pre-surgical EI values also showed a modest yet significant correlation to the ASRM scores calculated at the time of surgery ($R=0.6$, $p < 0.001$) and the cumulative distribution of EIs shows different and increasing mean values passing from control women to women with peritoneal, ovarian and deeply infiltrating endometriosis (Table 2: p for trend < 0.001) (Figure 2).

A Logistic regression model was used to estimate the risk to have DIE, adjusting for age of women. The analysis showed that women with EI score > 28 (75th percentile) have more than 24 times more risk to have DIE compared to women with EI score ≤ 28 (OR = 24.7, 95%CI = 8.3 to 73.7, $p < 0.001$).

DISCUSSION

Endometriosis cannot always be cured successfully by current medical and surgical interventions, and attempts for early diagnosis have been overwhelmed by “hit-and-miss” treatments (31). In particular, the lack of a non-invasive method of diagnosis has limited the ability of the clinician to choose appropriate treatment or surgery (25). In this study we have tested the clinical applications of a software-derived score, the Endometriosis Index, created to identify women affected by the disease.

The first step of the project has been the realization of an electronic aid able to save and elaborate the data derived by women at risk of bearing endometriosis; the fields requested by the software to be filled have been specifically designed for a multidisciplinary management of this disease (i.e. gynaecologists, anaesthetists, general surgeons, urologists, radiologists, pathologists etc.) and contain clinical information derived by the patient, the doctors and the diagnostic evidences (<http://www.galliera.it/endometriosi/promoie.html>). The first three panels collect data regarding symptoms and the level of patient disability. Dysmenorrhea, non-cyclic pelvic pain and dyspareunia are three distinct bars as VAS (Fig.1); words used to describe pain were brought together, categorized and scaled on a common intensity dimension, in accord to the McGill pain questionnaire (23, 32). Quality of life was determined by assessing the loss of productivity (days of work or study lost in the last month), the inability to perform daily life activities and sleep impairment (nights lost in the last month) (Fig.2). The gastrointestinal dysfunctions, urinary symptoms and headache were accurately listed and scaled in accord with recent literature (10, 11, 19, 26, 33) (Fig.3). The last panel contains and quantifies the positivity of the pelvic examination manoeuvres and diagnostic tests demonstrated to reveal the presence of endometriosis (3, 4, 7, 13, 21, 25, 34, 35).

At this point the challenge has been to find a way to convert all these clinical data into a number correlated to the presence and severity of endometriosis at the time of the consultation. In order to achieve this goal the clinical variables, their relative level of importance and the algorithm for the software were chosen after an exhaustive literature review and our previous practice (9-13, 17). In the final phase of the project we have validated this novel technology by testing in different situations: pre-surgical presence of endometriosis, absence of disease and post-excision endometriosis-free status.

The results of the present work show that women with endometriotic lesions had pre-operatively EI values higher than those calculated in patients with other benign pathologies (Tab.1 – Fig.1). The significant EI drops observed at consultation after the excision of endometriosis further exemplifies the capability of this non-invasive tool to reveal the presence/absence of disease. To our knowledge this is the first report of a dedicated software able to confirm the effect -- and effectiveness -- of surgical excision on endometriosis (Abbott, Howe et al. 2004) in patients managed for pain or impaired quality of life.

In addition, it is worth noting that pre-surgical EI values of patients affected by endometriosis correlate with the ASRM scores calculated at laparoscopy and their averages differ in the three forms of the disease. We consider it significant that the EI scores subgroup women affected by endometriosis in patients with peritoneal, ovarian and deeply infiltrating lesions (Table 2). The evidence that EI levels rise from ‘mild’ to ‘severe’ forms of endometriosis emphasises the sensitivity of the algorithm-regulated software and confirms the difficulty of predicting peritoneal instead of ovarian or DIE (25). At present, superficial endometriosis is considered a normal phenomenon in women at the childbearing age, whereas deep infiltrative endometriosis and endometrial ovarian cysts are the complex and painful manifestations of the condition (36). In accord to these latter evidences we must point out that high EI scores are strongly predictive of aggressive forms of endometriosis even though initial or mild types

of disease may be not diagnosed with certainty before surgery. The parameters/variables constitutive of the software, in fact, seem to better unmask the invasiveness of endometriosis rather than the disease itself. Gastrointestinal and genitourinal symptoms temporally related to the menstrual cycle are suggestive of retroperitoneal foci of disease and are more heavily weighted in the algorithm. These data could explain why the multivariate model showed that an EI score > 28 (75th percentile) is strongly associated (24-fold increased risk) to the presence of DIE compared to a lower EI score.

The data presented indicate the actual potential of a software-assisted clinical management of patients suspected to bear endometriosis. The first scenario is represented by patients with low EI scores in whom the empirical use of medical therapy could be justified before further consultations. On the contrary, women with alerting / non-diagnostic EI values should be carefully counselled for surgery considering both complications and clinical variables as age, course of infertility, quality of life etc. It is novel that an electronic tool could allow the selection of patients with a high score value to become candidates for a specific presurgical diagnostic work-up to confirm the presence of DIE. A correct preoperative diagnosis is of paramount importance in order to plan an adequate surgical procedure, and minimizes the risks of overlooking endometriotic intestinal or urinary lesions at surgery or of performing an incomplete surgical resection (15). This may eventually help to minimize the postoperative persistence or recurrence of both the lesions and the pain symptoms.

Although the number of patients can be considered still too low to allow definitive conclusions the pioneering use of the EI score as a screening tool to detect endometriosis proved clinically very effective and offered our patients the possibility to receive an early management of the disease, specific consents, and a dedicated surgical team.

The authors feel that this EI can serve as a non-invasive diagnostic tool, but, can also be useful in the endometriosis community as a standardized communication method. A large multi-centred randomized trial is required to validate this index.

Acknowledgments

The authors wish to thank as scientific adviser Dr. Mario Meroni of the Gynecological Unit of Niguarda Cà Granda Hospital in Milan.

References

1. Ballweg ML. Big picture of endometriosis helps provide guidance on approach to teens: comparative historical data show endo starting younger, is more severe. *J Pediatr Adolesc Gynecol* 2003;16:S21-6.
2. Wheeler JM. Epidemiology of endometriosis-associated infertility. *J Reprod Med* 1989;34:41-6.
3. Howard FM. Chronic pelvic pain. *Obstet Gynecol* 2003;101:594-611.
4. Gelbaya TA, El-Halwagy HE. Focus on primary care: chronic pelvic pain in women. *Obstet Gynecol Surv* 2001;56:757-64.
5. Hurd WW. Criteria that indicate endometriosis is the cause of chronic pelvic pain. *Obstet Gynecol* 1998;92:1029-32.
6. Solnik MJ. Chronic pelvic pain and endometriosis in adolescents. *Curr Opin Obstet Gynecol* 2006;18:511-8.
7. Garry R. Diagnosis of endometriosis and pelvic pain. *Fertil Steril* 2006;86:1307-9; discussion 17.
8. Abbott J, Hawe J, Hunter D, Holmes M, Finn P, Garry R. Laparoscopic excision of endometriosis: a randomized, placebo-controlled trial. *Fertil Steril* 2004;82:878-84.
9. Vercellini P, Fedele L, Aimi G, Pietropaolo G, Consonni D, Crosignani PG. Association between endometriosis stage, lesion type, patient characteristics and severity of pelvic pain symptoms: a multivariate analysis of over 1000 patients. *Hum Reprod* 2007;22:266-71.
10. Chapron C, Barakat H, Fritel X, Dubuisson JB, Breart G, Fauconnier A. Presurgical diagnosis of posterior deep infiltrating endometriosis based on a standardized questionnaire. *Hum Reprod* 2005;20:507-13.

11. Fauconnier A, Chapron C, Dubuisson JB, Vieira M, Dousset B, Breart G. Relation between pain symptoms and the anatomic location of deep infiltrating endometriosis. *Fertil Steril* 2002;78:719-26.
12. Davis AR, Westhoff CL. Primary dysmenorrhea in adolescent girls and treatment with oral contraceptives. *J Pediatr Adolesc Gynecol* 2001;14:3-8.
13. Porpora MG, Koninckx PR, Piazze J, Natili M, Colagrande S, Cosmi EV. Correlation between endometriosis and pelvic pain. *J Am Assoc Gynecol Laparosc* 1999;6:429-34.
14. Revised American Society for Reproductive Medicine classification of endometriosis: 1996. *Fertil Steril* 1997;67:817-21.
15. Chapron C, Dubuisson JB, Chopin N, Foulot H, Jacob S, Vieira M *et al.* [Deep pelvic endometriosis: management and proposal for a "surgical classification"]. *Gynecol Obstet Fertil* 2003;31:197-206.
16. Cornillie FJ, Oosterlynck D, Lauweryns JM, Koninckx PR. Deeply infiltrating pelvic endometriosis: histology and clinical significance. *Fertil Steril* 1990;53:978-83.
17. Reina SA, Reina VM, Debbia EA. Records matching model for data survey on applied and experimental microbiology. *New Microbiol* 2007;30:35-44.
18. Peveler R, Edwards J, Daddow J, Thomas E. Psychosocial factors and chronic pelvic pain: a comparison of women with endometriosis and with unexplained pain. *J Psychosom Res* 1996;40:305-15.
19. Ferrero S, Pretta S, Bertoldi S, Anserini P, Remorgida V, Del Sette M *et al.* Increased frequency of migraine among women with endometriosis. *Hum Reprod* 2004;19:2927-32.
20. Ferrero S, Esposito F, Abbamonte LH, Anserini P, Remorgida V, Ragni N. Quality of sex life in women with endometriosis and deep dyspareunia. *Fertil Steril* 2005;83:573-9.

21. Kennedy S, Bergqvist A, Chapron C, D'Hooghe T, Dunselman G, Greb R *et al.* ESHRE guideline for the diagnosis and treatment of endometriosis. *Hum Reprod* 2005;20:2698-704.
22. Pastor-Navarro H, Gimenez-Bachs JM, Donate-Moreno MJ, Pastor-Guzman JM, Ruiz-Mondejar R, Atienzar-Tobarra M *et al.* Update on the diagnosis and treatment of bladder endometriosis. *Int Urogynecol J Pelvic Floor Dysfunct* 2007;18:949-54.
23. Melzack R. The short-form McGill Pain Questionnaire. *Pain* 1987;30:191-7.
24. Jensen MP, Turner JA, Romano JM. Chronic pain coping measures: individual vs. composite scores. *Pain* 1992;51:273-80.
25. Eskenazi B, Warner M, Bonsignore L, Olive D, Samuels S, Vercellini P. Validation study of nonsurgical diagnosis of endometriosis. *Fertil Steril* 2001;76:929-35.
26. Elliott TE, Renier CM, Palcher JA. Chronic pain, depression, and quality of life: correlations and predictive value of the SF-36. *Pain Med* 2003;4:331-9.
27. Fauconnier A, Chapron C. Endometriosis and pelvic pain: epidemiological evidence of the relationship and implications. *Hum Reprod Update* 2005;11:595-606.
28. Jones G, Jenkinson C, Taylor N, Mills A, Kennedy S. Measuring quality of life in women with endometriosis: tests of data quality, score reliability, response rate and scaling assumptions of the Endometriosis Health Profile Questionnaire. *Hum Reprod* 2006;21:2686-93.
29. Fedele L, Bianchi S, Carmignani L, Berlanda N, Fontana E, Frontino G. Evaluation of a new questionnaire for the presurgical diagnosis of bladder endometriosis. *Hum Reprod* 2007;22:2698-701.
30. Dubernard G, Rouzier R, David-Montefiore E, Bazot M, Darai E. Use of the SF-36 questionnaire to predict quality-of-life improvement after laparoscopic colorectal resection for endometriosis. *Hum Reprod* 2008;23:846-51.

31. Bianconi L, Hummelshoj L, Coccia ME, Vigano P, Vittori G, Veit J *et al.* Recognizing endometriosis as a social disease: the European Union-encouraged Italian Senate approach. *Fertil Steril* 2007;88:1285-7.
32. Melzack R. The McGill pain questionnaire: from description to measurement. *Anesthesiology* 2005;103:199-202.
33. Chapron C, Chopin N, Borghese B, Foulot H, Dousset B, Vacher-Lavenu MC *et al.* Deeply infiltrating endometriosis: pathogenetic implications of the anatomical distribution. *Hum Reprod* 2006;21:1839-45.
34. Koninckx PR, Meuleman C, Oosterlynck D, Cornillie FJ. Diagnosis of deep endometriosis by clinical examination during menstruation and plasma CA-125 concentration. *Fertil Steril* 1996;65:280-7.
35. Biscaldi E, Ferrero S, Remorgida V, Rollandi GA. Bowel endometriosis: CT-enteroclysis. *Abdom Imaging* 2007;32:441-50.
36. De Ceglie A, Bilardi C, Bianchi S, Picasso M, Di Muzio M, Trimarchi A *et al.* Acute small bowel obstruction caused by endometriosis: A case report and review of the literature. *World J Gastroenterol* 2008;14:3430-4.

FIGURE LEGENDS

Figure 1. Boxplots of EI score before and after surgery in controls and in endometriosis categories.

Figure 2. Panels of IE-Ring Software: **A)** Pain assessment is the result of the following subitems: *1 Dysmenorrhea (IDM) + 2 Non menstrual pelvic pain (IDP) + 3 Dyspareunia (DRS)*. **B)** Characterization of the sensory dimension of pain (CDD) is the result of the following subitems: *4 Throbbing + 5 Stabbing + 6 Cramping + 7 Hot-burning + 8 Heavy + 9 Tender*. Quality of life limitation (LDD) is the result of the following subitems: *10 Work/School days of absence + 11 Daily activity restriction and 12 Sleep impairment*. **C)** Induced dysfunction and physical alteration (SPA) is the result of the following subitems: gastrointestinal symptoms (*13 Alternating constipation and diarrhoea + 14 Rectal tenesmus + 15 constipation + 16 diarrhoea + 17 Rectal pain*), urinary symptoms (*18 Urinary pain – 19 bladder tenesmus – 20 Frequent urination – 21 Dysuria*) and headache (*22 Headache – 23 Migraine – 24 cluster headache*). **D)** Pelvic examination (VM) is the result of the following subitems: (*25 Grade of fixed uterus + 26 Presence of adnexal cyst(s) + 27 Grade of Douglas tenderness + 28 Presence of infiltrating nodule(s) + 29 Grade of pain at digital exploration*); Diagnostic evidences (IS) result of the following subitems: (*30 CA-125 blood assay + 31 Ultrasound finding(s) + 32 CA 19.9 blood assay + 33 Magnetic resonance finding(s) + 34 X-Ray / Computerized Tomography finding(s) + 35 Colonoscopy finding(s)*).